

## Curve Sketching

Example.  $f(x) = \frac{2x^2}{x^2 - 1}$

A. Domain.  $f(x)$  is undefined when  $x^2 - 1 = 0 \Leftrightarrow (x - 1)(x + 1) = 0 \Leftrightarrow x = 1, -1$ .  
Thus  $\text{Domain}(f) = (-\infty, -1) \cup (-1, 1) \cup (1, +\infty)$ .

B. Intercepts.  $y$ -intercept:  $y(0) = 0$ .

$x$ -intercepts:  $f(x) = 0 \Leftrightarrow \frac{2x^2}{x^2 - 1} = 0 \Leftrightarrow x = 0$ .

Thus  $(0, 0)$  is the only intercept.

C. Symmetry.  $\frac{2(-x)^2}{(-x)^2 - 1} = \frac{2x^2}{x^2 - 1} = 0$

i.e.  $f(-x) = f(x)$ , i.e.  $f(x)$  is even, therefore the graph is symmetric about the  $y$ -axis.

D. Asymptotes.

horizontal:  $\lim_{x \rightarrow +\infty} \frac{2x^2}{x^2 - 1} = \lim_{x \rightarrow +\infty} \frac{2}{1 - \frac{1}{x^2}} = 2 \Rightarrow y = 2$  is a horizontal asymptote.

$$\lim_{x \rightarrow -\infty} \frac{2x^2}{x^2 - 1} = \lim_{x \rightarrow -\infty} \frac{2}{1 - \frac{1}{x^2}} = 2$$

vertical:  $\lim_{x \rightarrow 1^+} \frac{2x^2}{x^2 - 1} \left( = \frac{2}{+0} \right) = +\infty$

$$\lim_{x \rightarrow 1^-} \frac{2x^2}{x^2 - 1} \left( = \frac{2}{-0} \right) = -\infty$$

$\Rightarrow x = 1$  and  $x = -1$  are vertical asymptotes.

$$\lim_{x \rightarrow -1^+} \frac{2x^2}{x^2 - 1} \left( = \frac{2}{-0} \right) = -\infty$$

$$\lim_{x \rightarrow -1^-} \frac{2x^2}{x^2 - 1} \left( = \frac{2}{+0} \right) = +\infty$$

E. Increase / decrease.

$$f'(x) = \frac{4x(x^2 - 1) - 2x^2 \cdot 2x}{(x^2 - 1)^2} = \frac{-4x}{(x^2 - 1)^2}$$

since  $(x^2 - 1)^2 > 0$  for  $x \neq \pm 1$ ,

$f'(x) > 0$  when  $x < 0 \Rightarrow f(x)$  is increasing on  $(-\infty, -1)$  and on  $(-1, 0)$ .

$f'(x) < 0$  when  $x > 0 \Rightarrow f(x)$  is decreasing on  $(0, 1)$  and on  $(1, +\infty)$ .

F. Max / min.

$$f'(x) = 0 \Rightarrow \frac{2x^2}{x^2 - 1} = 0 \Rightarrow x = 0.$$

At 0,  $f'(x)$  changes from + to -, so 0 is a local maximum.

$f'(x)$  does not exist when  $x^2 - 1 = 0 \Rightarrow x = \pm 1$ , but 1 and -1 are not in the domain of  $f(x)$ , so they are not critical numbers.

G. Concavity and inflection points.

$$f''(x) = \frac{-4(x^2 - 1)^2 - (-4x)2(x^2 - 1)2x}{(x^2 - 1)^4} = \frac{(x^2 - 1)[-4(x^2 - 1) + 4x \cdot 2 \cdot 2x]}{(x^2 - 1)^4} =$$

$$= \frac{-4x^2 + 4 + 16x^2}{(x^2 - 1)^3} = \frac{12x^2 + 4}{(x^2 - 1)^3}.$$

since  $12x^2 + 4 > 0$  for all  $x$ ,

$f''(x) > 0$  when  $x^2 - 1 > 0 \Rightarrow f(x)$  is CU on  $(-\infty, -1)$  and on  $(1, +\infty)$ .

$f''(x) < 0$  when  $x^2 - 1 < 0 \Rightarrow f(x)$  is CD on  $(-1, 1)$ .


$f(x)$  changes the direction of concavity at  $\pm 1$ , but  $1$  and  $-1$  are not in the domain of  $f(x)$ , so there are no inflection points.

H. Graph.

first sketch the asymptotes (also notice that when  $x$  is large or large negative,

$f(x) = \frac{2x^2}{x^2 - 1} = \frac{2}{1 - \frac{1}{x^2}}$  is slightly bigger than 2 because  $1 - \frac{1}{x^2}$  is slightly smaller than 1,

so the graph approaches the horizontal asymptote from above), plot the intercepts (only  $(0, 0)$  in our case), maxima (none), minima (none), and inflection points (none), and indicate the intervals of increase, decrease, and concavity:



now sketch the graph:

